

## University of Groningen

### Air Pollution and Respiratory Infections during Early Childhood

MacIntyre, Elaina A.; Gehring, Ulrike; Moelter, Anna; Fuertes, Elaine; Kluemper, Claudia; Kraemer, Ursula; Quass, Ulrich; Hoffmann, Barbara; Gascon, Mireia; Brunekreef, Bert

*Published in:*  
Environmental Health Perspectives

*DOI:*  
[10.1289/ehp.1306755](https://doi.org/10.1289/ehp.1306755)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2014

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

MacIntyre, E. A., Gehring, U., Moelter, A., Fuertes, E., Kluemper, C., Kraemer, U., Quass, U., Hoffmann, B., Gascon, M., Brunekreef, B., Koppelman, G. H., Beelen, R., Hoek, G., Birk, M., de Jongste, J. C., Smit, H. A., Cyrus, J., Gruziova, O., Korek, M., ... Heinrich, J. (2014). Air Pollution and Respiratory Infections during Early Childhood: An Analysis of 10 European Birth Cohorts within the ESCAPE Project. *Environmental Health Perspectives*, 122(1), 107-113. <https://doi.org/10.1289/ehp.1306755>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Air Pollution and Respiratory Infections during Early Childhood: An Analysis of 10 European Birth Cohorts within the ESCAPE Project

Elaina A. MacIntyre,<sup>1</sup> Ulrike Gehring,<sup>2</sup> Anna Mölter,<sup>3</sup> Elaine Fuertes,<sup>1,4</sup> Claudia Klümper,<sup>5</sup> Ursula Krämer,<sup>5</sup> Ulrich Quass,<sup>6</sup> Barbara Hoffmann,<sup>5,7</sup> Mireia Gascon,<sup>8,9</sup> Bert Brunekreef,<sup>2,10</sup> Gerard H. Koppelman,<sup>11,12</sup> Rob Beelen,<sup>2</sup> Gerard Hoek,<sup>2</sup> Matthias Birk,<sup>1</sup> Johan C. de Jongste,<sup>13</sup> H.A. Smit,<sup>10</sup> Josef Cyrys,<sup>14</sup> Olena Gruzieva,<sup>15</sup> Michal Korek,<sup>15</sup> Anna Bergström,<sup>15</sup> Raymond M. Agius,<sup>3</sup> Frank de Vocht,<sup>3</sup> Angela Simpson,<sup>16</sup> Daniela Porta,<sup>17</sup> Francesco Forastiere,<sup>17</sup> Chiara Badaloni,<sup>17</sup> Giulia Cesaroni,<sup>17</sup> Ana Esplugues,<sup>9,18</sup> Ana Fernández-Somoano,<sup>9,19</sup> Aitana Lerxundi,<sup>20,21</sup> Jordi Sunyer,<sup>8,9,22,23</sup> Marta Cirach,<sup>8,9</sup> Mark J. Nieuwenhuijsen,<sup>8,9</sup> Göran Pershagen,<sup>15</sup> and Joachim Heinrich<sup>1</sup>

<sup>1</sup>Institute of Epidemiology I, Helmholtz Zentrum München, German Research Centre for Environmental Health, Munich, Germany; <sup>2</sup>Institute for Risk Assessment Sciences, Utrecht University, Utrecht, the Netherlands; <sup>3</sup>Centre for Epidemiology, Institute of Population Health, Manchester Academic Health Sciences Centre, The University of Manchester, Manchester, United Kingdom; <sup>4</sup>School of Population and Public Health, University of British Columbia, Vancouver, British Columbia, Canada; <sup>5</sup>IUF-Leibniz Research Institute for Environmental Medicine, University of Düsseldorf, Düsseldorf, Germany; <sup>6</sup>Air Quality & Sustainable Nanotechnology, IUTA (Institut für Energie-und Umwelttechnik e.V.), Duisburg, Germany; <sup>7</sup>Medical Faculty, Heinrich Heine University Düsseldorf, Düsseldorf, Germany; <sup>8</sup>Centre for Research in Environmental Epidemiology (CREAL), Barcelona, Catalonia, Spain; <sup>9</sup>Spanish Consortium for Research on Epidemiology and Public Health (CIBERESP), Spain; <sup>10</sup>Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, the Netherlands; <sup>11</sup>Department of Pediatric Pulmonology and Pediatric Allergology, Beatrix Children's Hospital, and <sup>12</sup>GRIAC (Groningen Research Institute of Asthma and COPD), University of Groningen, University Medical Center Groningen, Groningen, the Netherlands; <sup>13</sup>Department of Pediatrics, Division of Respiratory Medicine, Erasmus University Medical Center/Sophia Children's Hospital, Rotterdam, the Netherlands; <sup>14</sup>Institute of Epidemiology II, Helmholtz Zentrum München, German Research Centre for Environmental Health, Munich, Germany; <sup>15</sup>Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden; <sup>16</sup>Institute of Inflammation and Repair, Manchester Academic Health Sciences Centre, The University of Manchester, Manchester, United Kingdom; <sup>17</sup>Department of Epidemiology, Lazio Regional Health Service ASL RME (Dipartimento di epidemiologia ASL Roma E), Rome, Italy; <sup>18</sup>Centre of Public Health Research (CSISP), Valencia, Spain; <sup>19</sup>Department of Preventive Medicine, University of Oviedo, Asturias, Spain; <sup>20</sup>BioDonostia Health Research Institute, Donostia, Spain; <sup>21</sup>Department of Public Health and Preventive Medicine, University of Basque Country, Bilbao, Spain; <sup>22</sup>Hospital del Mar Medical Research Institute (IMIM), Barcelona, Catalonia, Spain; <sup>23</sup>Pompeu Fabra University, Department of Experimental and Health Sciences, Barcelona, Catalonia, Spain

**BACKGROUND:** Few studies have investigated traffic-related air pollution as a risk factor for respiratory infections during early childhood.

**OBJECTIVES:** We aimed to investigate the association between air pollution and pneumonia, croup, and otitis media in 10 European birth cohorts—BAMSE (Sweden), GASPII (Italy), GINIplus and LISAPLUS (Germany), MAAS (United Kingdom), PIAMA (the Netherlands), and four INMA cohorts (Spain)—and to derive combined effect estimates using meta-analysis.

**METHODS:** Parent report of physician-diagnosed pneumonia, otitis media, and croup during early childhood were assessed in relation to annual average pollutant levels [nitrogen dioxide (NO<sub>2</sub>), nitrogen oxide (NO<sub>x</sub>), particulate matter ≤ 2.5 μm (PM<sub>2.5</sub>), PM<sub>2.5</sub> absorbance, PM<sub>10</sub>, PM<sub>2.5-10</sub> (coarse PM)], which were estimated using land use regression models and assigned to children based on their residential address at birth. Identical protocols were used to develop regression models for each study area as part of the ESCAPE project. Logistic regression was used to calculate adjusted effect estimates for each study, and random-effects meta-analysis was used to calculate combined estimates.

**RESULTS:** For pneumonia, combined adjusted odds ratios (ORs) were elevated and statistically significant for all pollutants except PM<sub>2.5</sub> (e.g., OR = 1.30; 95% CI: 1.02, 1.65 per 10-μg/m<sup>3</sup> increase in NO<sub>2</sub> and OR = 1.76; 95% CI: 1.00, 3.09 per 10-μg/m<sup>3</sup> PM<sub>10</sub>). For otitis media and croup, results were generally null across all analyses except for NO<sub>2</sub> and otitis media (OR = 1.09; 95% CI: 1.02, 1.16 per 10-μg/m<sup>3</sup>).

**CONCLUSION:** Our meta-analysis of 10 European birth cohorts within the ESCAPE project found consistent evidence for an association between air pollution and pneumonia in early childhood, and some evidence for an association with otitis media.

**CITATION:** MacIntyre EA, Gehring U, Mölter A, Fuertes E, Klümper C, Krämer U, Quass U, Hoffmann B, Gascon M, Brunekreef B, Koppelman GH, Beelen R, Hoek G, Birk M, de Jongste JC, Smit HA, Cyrys J, Gruzieva O, Korek M, Bergström A, Agius RM, de Vocht F, Simpson A, Porta D, Forastiere F, Badaloni C, Cesaroni G, Esplugues A, Fernández-Somoano A, Lerxundi A, Sunyer J, Cirach M, Nieuwenhuijsen MJ, Pershagen G, Heinrich J. 2014. Air pollution and respiratory infections during early childhood: an analysis of 10 European birth cohorts within the ESCAPE project. *Environ Health Perspect* 122:107–113; <http://dx.doi.org/10.1289/ehp.1306755>

## Introduction

Respiratory infections are a leading reason for outpatient physician visits and hospitalizations among children (Williams et al. 2002). Most infections resolve with minimal use of health care resources; however, episodes of severe or recurrent infection may require hospitalization or surgery, and the resultant burden on resources is substantial (Black et al. 2010).

Young children are particularly susceptible to respiratory pathogens and also to air pollution (Bateson and Schwartz 2008;

Heinrich and Slama 2007). There is strong evidence that indoor air pollution, such as secondhand smoke and the use of biomass, is a risk factor for respiratory infections in children (da Costa et al. 2004). Evidence is growing to support an association with outdoor air pollution as well (Brauer et al. 2006; Leonardi et al. 2000; MacIntyre et al. 2011).

The European Study of Cohorts for Air Pollution Effects (ESCAPE 2009) is a project aimed at investigating the impacts of long-term exposure to air pollution through the development of harmonized exposure

data assigned to previously established cohorts that have collected information on specific health outcomes of interest for air pollution research. We analyzed data from

Address correspondence to J. Heinrich, Institute of Epidemiology I, Helmholtz Zentrum München, Ingolstädter Landstraße 1, 85764 Neuherberg Germany. Telephone: 49-(0)89-3187-4150. E-mail: [heinrich@helmholtz-muenchen.de](mailto:heinrich@helmholtz-muenchen.de)

Supplemental Material is available online (<http://dx.doi.org/10.1289/ehp.1306755>).

The research leading to these results was funded by the European Community's Seventh Framework Program (FP7/2007–2011) under grant 211250. The BAMSE study was supported by the Swedish Research Council FORMAS (for Environment, Agricultural Sciences and Spatial Planning), the Stockholm County Council, the Swedish Foundation for Health Care Sciences and Allergy Research, and the Swedish Environmental Protection Agency. The GINIplus study was supported for the first 3 years by the Federal Ministry for Education, Science, Research and Technology, Germany (interventional arm) and Helmholtz Zentrum München, Germany (former GSF; National Research Center for Environment and Health) (observational arm). The LISAPLUS study was supported by grants from the Federal Ministry for Education, Science, Research and Technology, Germany; Helmholtz Zentrum München, Germany (former GSF); Helmholtz Centre for Environmental Research—UFZ, Germany; Marien-Hospital Wesel, Germany; and Pediatric Practice, Bad Honnef, Germany. The PIAMA study is supported by The Netherlands Organization for Health Research and Development; The Netherlands Organization for Scientific Research; The Netherlands Asthma Fund; The Netherlands Ministry of Spatial Planning, Housing, and the Environment; and The Netherlands Ministry of Health, Welfare, and Sport. MAAS was supported by an Asthma UK Grant (04/014); the JP Moulton Charitable Foundation, UK; and the James Trust and Medical Research Council, UK (G0601361). INMA was funded by grants from the Spanish Ministry of Health-Instituto de Salud Carlos III (Red INMA G03/176, CB06/02/0041, FIS-PI041436, FIS-PI081151, FIS-PI042018, FIS-PI09/02311, FIS-PI06/0867, FIS-PS09/00090, FIS-FEDER 03/1615, 04/1509, 04/1112, 04/1931, 05/1079, 05/1052, 06/1213, 07/0314, and 09/02647); Generalitat de Catalunya-CIRIT, Spain (1999SGR 00241); Conselleria de Sanitat Generalitat Valenciana, Spain; Universidad de Oviedo, Obra social Cajastur, Spain; Department of Health of the Basque Government, Spain (2005111093 and 2009111069); Provincial Government of Gipuzkoa (DFG06/004 and DFG08/001), Spain; and Fundación Roger Torné, Spain. GASPII was funded by The Italian Ministry of Health (ex art.12 D.Lgs 502/92, 2001).

The authors declare they have no actual or potential competing financial interests.

Received: 4 March 2013; Accepted: 30 September 2013; Advance Publication: 22 October 2013; Final Publication: 1 January 2014.

10 European birth cohorts and completed a meta-analysis of air pollution and respiratory infection (pneumonia, croup, and otitis media) during early childhood.

## Methods

**Study population.** We included 10 ESCAPE birth cohorts. The inclusion criteria for each birth cohort were that data on at least one outcome of interest were available during early childhood, and that the ESCAPE exposure assignment was complete.

BAMSE (Children, Allergy, Milieu, Stockholm, Epidemiological Survey) is a population-based prospective birth cohort of children born during 1994–1996 in Stockholm County, Sweden (Wickman et al. 2002). GASPII (Gene and Environment: Prospective Study on Infancy in Italy) is a prospective birth cohort of children born during 2003–2004 in Rome, Italy (Porta et al. 2007). GINIplus (German Infant Nutrition Intervention Study Plus environmental and genetic influences on allergy development) is a population-based prospective birth cohort, with a nutritional intervention, of children born during 1995–1998 in Wesel and Munich, Germany (Zirngibl et al. 2002). LISAPLUS (Influence of Life-style Factors on the Development of the Immune System and Allergies in Childhood Plus the influence of traffic emissions and genetics) is a population-based prospective birth cohort study of children born during 1997–1999 in Wesel, Munich, Leipzig, and Bad Honnef, Germany (Heinrich et al. 2002). INMA (Infancia y Medio Ambiente; Environment and Childhood) is a network of Spanish birth cohorts. The four INMA cohorts in the present analysis comprise children born during 2004–2008 in both major cities and rural towns—Asturias, Gipuzkoa, Sabadell, and Valencia (Guxens et al. 2012). MAAS (Manchester Asthma and Allergy Study) is an unselected, prospective population-based birth cohort study (with a small nested allergen control intervention) of children born during 1995–1997 in the Greater Manchester conurbation in the United Kingdom (Custovic et al. 2002). Finally, the PIAMA (Prevention and Incidence of Asthma and Mite Allergy) study is a population based prospective birth cohort, with an intervention component, of children born during 1996–1997 in cities and small towns across the Netherlands (Brunekreef et al. 2002). Each cohort obtained parental consent and protocols were approved by all relevant ethical review boards.

Exposure to outdoor air pollution was estimated using land use regression (LUR) models, and methods were harmonized across each cohort (Beelen et al. 2013; Cyrus et al. 2012; Eeftens et al. 2012a, 2012b; ESCAPE 2013). Briefly, sampling sites for particulate

matter ( $n = 20$ – $40$ ) and nitrogen oxides ( $n = 40$ – $80$ ) were strategically chosen for each study area to represent the spatial distribution of concentrations at the residential address of each child at birth, with some overrepresentation of locations with heavy traffic where the largest heterogeneity was expected. Three 2-week sampling campaigns were spread out over 1 year and used to estimate an annual average. Measurement data used to develop the LUR models were collected during 2008–2009 (BAMSE, GINIplus/LISAPLUS, PIAMA), 2009 (INMA), 2009–2010 (MAAS), and 2010–2011 (GASPII). The annual average was temporally adjusted using continuous measurement data from a fixed monitor that was used to capture background levels in each study area. Models based on GIS (geographic information system) variables related to traffic, land use, population density, altitude, and regional background pollution were developed using common ESCAPE protocols to predict measured air pollution concentrations.

Separate models were developed for nitrogen dioxide ( $\text{NO}_2$ ), nitrogen oxides [ $\text{NO}_x$  ( $\text{NO}_2 + \text{NO}$ )],  $\text{PM}_{2.5}$  (particulate matter  $\leq 2.5 \mu\text{m}$  in diameter),  $\text{PM}_{2.5}$  absorbance,  $\text{PM}_{10}$  ( $\leq 10 \mu\text{m}$  in diameter), and coarse PM ( $\text{PM}_{10}$ – $\text{PM}_{2.5}$ ). In addition, two variables were created to describe traffic intensity at the residential address: traffic intensity on the nearest street, and traffic load on all major roads within a 100-m buffer. Annual average air pollution concentrations and traffic intensity variables were assigned to children for the first year of life based on their residential address reported at birth. The LISAPLUS study centers of Leipzig and Bad Honnef were not included in the ESCAPE exposure assessment, so children from these cities could not be included in the meta-analysis.

The air pollution data used to derive the ESCAPE exposure models were measured in 2008–2011, whereas children included in the study cohorts were born as early as 1994. Therefore, we conducted sensitivity analyses using routine monitoring data to back-extrapolate exposure estimates based on LUR to each child's year of birth. We used two approaches for the back-extrapolation: The first used the ratio of the average concentration measured from the date of birth through the second birthday to the average concentration measured during the ESCAPE monitoring year; the second used the absolute difference between the average concentrations at each time period (ESCAPE 2013). Both methods altered the spatial contrast derived from the current LUR models without affecting the spatial patterns of air pollutants in the study areas.

For each cohort, parents reported (yes/no) physician-diagnosed pneumonia, otitis media, and croup during early childhood (see

Supplemental Material, p. 2, for the specific questions used for each cohort). Outcomes were assessed at 6 months (GASPII, LISAPLUS), 1 year (BAMSE, GINIplus, INMA Valencia, LISAPLUS, PIAMA), 14 months (INMA Gipuzkoa, INMA Sabadell), 15 months (GASPII), 18 months (LISAPLUS, INMA Asturias), 2 years (BAMSE, GINIplus, LISAPLUS, PIAMA), and 3 years (MAAS). It was not possible to evaluate respiratory infections restricted to the first 2 years of life for the MAAS birth cohort because these outcomes were not assessed in the full cohort until 3 years of age. Pneumonia data were available for all cohorts; otitis media data were available for all except GINIplus and MAAS; and croup data were available for all except GASPII, the four INMA cohorts, and PIAMA. Cumulative incidence was modeled in each analysis, unless otherwise specified.

We used logistic binomial regression in all individual cohort analyses, and statistical significance was defined by  $p$ -values  $< 0.05$ . Air pollution was entered as a continuous variable and was not transformed. Models were assessed using the Hosmer–Lemeshow goodness-of-fit test and the Pearson's chi-square test. Potential confounders were identified from previous literature and selected *a priori*. Individual cohort models were adjusted for municipality/city (BAMSE only), sex, older siblings (any/none), partial or exclusive breastfeeding at 6 months, atopy of either parent, child-care attendance reported at any time during follow-up, maternal smoking during pregnancy, secondhand smoke in the home reported at any time during follow-up (not available for INMA), visible mold or dampness in the home, use of gas stove, birth season (winter: January–March; spring: April–June; summer: July–September; fall: October–December), parental socioeconomic status [highest education attained by either parent (BAMSE, GINIplus, LISAPLUS, PIAMA, INMA: low, medium, high); highest occupational level by either parent (GASPII: low, medium, high); or household income (MAAS:  $< \text{£}10,000$ ;  $\text{£}10,000$ – $20,000$ ;  $\text{£}20,000$ – $30,000$ ;  $> \text{£}30,000$ )], and intervention (GINIplus, MAAS, and PIAMA only). Models for traffic intensity and traffic load were additionally adjusted for background  $\text{NO}_2$  concentrations. Children with missing data for any covariate were excluded from individual analyses. Based on the ESCAPE protocol, we calculated estimates for the following increments in exposure:  $10 \mu\text{g}/\text{m}^3$  for  $\text{NO}_2$ ,  $20 \mu\text{g}/\text{m}^3$  for  $\text{NO}_x$ , 1 unit for  $\text{PM}_{2.5}$  absorbance,  $5 \mu\text{g}/\text{m}^3$  for  $\text{PM}_{2.5}$ ,  $10 \mu\text{g}/\text{m}^3$  for  $\text{PM}_{10}$ ,  $5 \mu\text{g}/\text{m}^3$  for coarse PM, 5,000 vehicles/day for traffic intensity on the nearest street; and 4,000 vehicle-km/day for traffic load on major roads within a 100-m buffer. We assessed heterogeneity of effect estimates



between studies using the  $I^2$  statistic (Huedo-Medina et al. 2006). We used random-effects meta-analysis models to calculate combined estimates (DerSimonian and Laird 1986).

We used sensitivity analyses to test the robustness of effect estimates to the inclusion of additional potential confounders: birth weight, maternal age at birth, and area-level socioeconomic indicators. In addition, we stratified associations for outcomes that were diagnosed during the first year of life and outcomes diagnosed during the second year of life for cohorts that completed follow-ups at 1 and 2 years of age (BAMSE, GINIplus, LISAPLUS, PIAMA). Additional analyses were stratified by sex, parental socioeconomic status (low, middle, or high), and residential mobility (moved from the birth address at any time during the follow-up period) to examine potential effect modification. As noted above, we also performed sensitivity analyses using exposure estimates that were recalculated for selected pollutants using back-extrapolation techniques to assess the consistency of associations. In addition, we performed a sensitivity analysis of the influence of neighborhood clustering by including an area-level variable (BAMSE: neighborhood; GINIplus: ZIP code; LISAPLUS: ZIP code; INMA: rural indicator; PIAMA: neighborhood) as a random effect in adjusted models. Area-level data were not available for GASPII or MAAS. Finally, we used two-pollutant models to estimate the independent effects of  $\text{NO}_2$  and PM.

All individual and combined analyses were completed using identical protocols. Individual estimates are presented by cohort except for the German birth cohorts (LISAPLUS and GINIplus), which had

almost identical study designs and parental questionnaires, and are presented as GINI/LISA North (Wesel) and GINI/LISA South (Munich) because separate air pollution models were developed for each area as part of ESCAPE. Statistical analyses were completed using SPSS version 20 (IBM SPSS, Armonk, NY, USA) and SAS version 9.1 (SAS Institute Inc., Cary NC, USA).

## Results

There was complete outcome (at least one), exposure (a minimum of  $\text{NO}_2$  and  $\text{NO}_x$ ) and potential confounder information for 16,059 children across all 10 cohorts (79.6% of the total recruited population). Children excluded due to missing data were more likely to have parents of lower socioeconomic status (BAMSE, GINI/LISA South, GINI/LISA North, MAAS, PIAMA), mothers who smoked during pregnancy (BAMSE, GASPII, GINI/LISA South, GINI/LISA North, PIAMA); and were less likely to be breastfed for at least 6 months (GASPII, GINI/LISA South, GINI/LISA North, MAAS, PIAMA) or to have atopic parents (GINI/LISA South, INMA Sabadell). Table 1 shows the cumulative incidence of parent-reported physician-diagnosed respiratory infection, by cohort. The cumulative incidence of pneumonia during early childhood ranged from 1.5% in INMA Sabadell to 7.9% in BAMSE (0.7–3.6% during the first year only). Otitis media ranged from 21.8% in GASPII to 50.0% in BAMSE (6.8–26.6% for the first year), and croup ranged from 10.6% in MAAS to 12.9% in GINI/LISA North (4.2–5.6% for the first year). There were differences in breastfeeding, child-care attendance, parental atopy, and secondhand smoke exposure

among the cohorts (see Supplemental Material, Table S1). Air pollution concentrations were highest in GASPII and lowest in BAMSE; GINI/LISA South, GINI/LISA North and PIAMA had similar mean concentrations (Table 1). Additional statistics on air pollutant concentrations by cohort are available in Supplemental Material, Table S2. Air pollutant concentrations were moderately to highly correlated (see Supplemental Material, Table S3; e.g., correlation between  $\text{PM}_{2.5}$  and  $\text{NO}_2$  ranged between 0.42 and 0.80, and correlations between  $\text{PM}_{2.5}$  absorbance and  $\text{NO}_2$  ranged between 0.40 and 0.93).

Associations between air pollution and respiratory infection during early childhood are presented in Figure 1 for *a*) individual and *b*) combined effect estimates. Table 2 presents combined effect estimates for crude (adjusted for sex and municipality) and adjusted (adjusted for all potential confounders) models and *p*-values for heterogeneity. The heterogeneity between studies varied and the largest  $I^2$  statistics were for models of pneumonia and  $\text{NO}_2$ ,  $\text{PM}_{2.5}$ , and  $\text{PM}_{10}$ . Effect estimates were robust to adjustment for older siblings, breastfeeding, parental atopy, child care, maternal smoking during pregnancy, environmental tobacco smoke, visible mold or dampness, use of gas stove, birth season, and parental socioeconomic status. For pneumonia, elevated odds ratios (ORs) were found in almost all analyses, and the combined estimates were statistically significant for all measures of air pollution except  $\text{PM}_{2.5}$  (OR = 2.58; 95% CI: 0.91, 7.27 for a  $5\text{-}\mu\text{g}/\text{m}^3$  increase). For otitis media and croup, results were generally null across all analyses except for  $\text{NO}_2$  and otitis media, for which the adjusted OR was 1.09 (95% CI: 1.02, 1.16 for a  $10\text{-}\mu\text{g}/\text{m}^3$  increase).

**Table 1.** The cumulative incidence of respiratory infections and distribution of air pollution for each ESCAPE birth cohort.

	BAMSE	GASPII	GINI/LISA South	GINI/LISA North	INMA Asturias	INMA Gipuzkoa	INMA Sabadell	INMA Valencia	MAAS	PIAMA
<b>Respiratory infections [n(%)]</b>										
Total included	3,821 (100)	678 (100)	3,321 (100)	2,460 (100)	360 (100)	437 (100)	402 (100)	559 (100)	695 (100)	3,475 (100)
Pneumonia, 0–1 year	137 (3.6)	5 (0.7) <sup>a</sup>	81 (2.4)	80 (3.3)	—	—	—	10 (1.8) <sup>b</sup>	—	84 (2.4)
Pneumonia, 0–2 years	301 (7.9)	14 (2.1) <sup>c</sup>	198 (6.0)	144 (5.9)	8 (2.2) <sup>d</sup>	9 (2.1) <sup>e</sup>	6 (1.5) <sup>e</sup>	—	13 (1.9) <sup>f</sup>	150 (4.3)
Otitis media, 0–1 year	1,017 (26.6) <sup>g</sup>	46 (6.8) <sup>g</sup>	202 (6.4) <sup>h</sup>	49 (18.2) <sup>h</sup>	—	—	—	139 (24.9) <sup>b</sup>	—	603 (18.0)
Otitis media, 0–2 years	1,911 (50.0) <sup>g</sup>	148 (21.8) <sup>c</sup>	422 (34.2) <sup>h</sup>	103 (38.3) <sup>h</sup>	143 (39.7) <sup>d</sup>	161 (36.8) <sup>e</sup>	144 (35.8) <sup>e</sup>	—	—	1,144 (32.9)
Croup, 0–1 year	201 (5.3)	—	140 (4.2)	135 (5.6)	—	—	—	—	—	—
Croup, 0–2 years	410 (10.7)	—	362 (10.9)	318 (12.9)	—	—	—	—	74 (10.6) <sup>f</sup>	—
<b>Air pollution (median [IQR])</b>										
$\text{NO}_2$ ( $\mu\text{g}/\text{m}^3$ )	12.4 (9.3)	43.2 (10.4)	20.8 (8.3)	23.2 (3.2)	22.2 (14.4)	18.4 (5.7)	41.7 (12.4)	27.9 (18.1)	23.0 (2.4)	23.1 (8.4)
$\text{NO}_x$ ( $\mu\text{g}/\text{m}^3$ )	20.9 (18.2)	65.8 (23.9)	34.6 (12.2)	33.3 (8.4)	44.0 (39.0)	37.2 (11.1)	69.7 (21.2)	44.7 (31.1)	38.7 (5.2)	32.8 (11.0)
$\text{PM}_{2.5}$ ( $\mu\text{g}/\text{m}^3$ )	8.1 (1.9)	18.8 (2.0)	13.3 (1.2)	17.2 (0.9)	—	—	14.6 (1.1)	—	9.4 (0.0)	16.5 (1.2)
$\text{PM}_{2.5}$ absorbance ( $10^{-5}/\text{m}$ )	0.6 (0.3)	2.5 (0.4)	1.7 (0.2)	1.2 (0.2)	—	—	2.2 (0.5)	—	1.1 (0.2)	1.2 (0.3)
$\text{PM}_{10}$ ( $\mu\text{g}/\text{m}^3$ )	15.6 (3.9)	34.9 (6.1)	20.4 (2.9)	25.2 (1.6)	—	—	26.4 (3.4)	—	17.0 (0.2)	24.6 (1.2)
Coarse PM ( $\mu\text{g}/\text{m}^3$ )	7.7 (3.0)	15.7 (4.3)	6.5 (2.0)	8.4 (0.7)	—	—	11.3 (2.5)	—	6.9 (0.8)	8.1 (0.8)
Traffic intensity on nearest street (vehicles/day)	500 (1,450)	—	500 (0)	500 (0)	—	—	—	—	500 (0)	215 (436)
Traffic load on major streets within 100-m buffer (vehicle $\times$ m/day)	0 (1,621,333)	—	0 (1,318,179)	0 (0)	—	—	—	—	0 (0)	0 (0)

IQR, interquartile range.

<sup>a</sup>Assessed at 6 months. <sup>b</sup>Assessed at 12 months. <sup>c</sup>Assessed at 15 months. <sup>d</sup>Assessed at 18 months. <sup>e</sup>Assessed at 14 months. <sup>f</sup>Assessed at 3 years. <sup>g</sup>Requiring antibiotic. <sup>h</sup>Otitis media was collected only in the LISAPLUS study (total South  $n = 1,242$ ; North  $n = 280$ ).

Effect estimates in two-pollutant models that included NO<sub>2</sub> plus one of the PM exposures were closer to the null (vs. estimates from single-pollutant models), and the only statistically significant finding was for NO<sub>2</sub> and otitis media (OR = 1.13; 95% CI: 1.01, 1.26 for a 10-μg/m<sup>3</sup> increase in NO<sub>2</sub>) when adjusted for coarse PM (see Supplemental Material, Table S4). Confidence intervals increased substantially in two-pollutant models, reflecting the high correlation between pollutants (see Supplemental Material, Table S3).

All measures of air pollution were associated with pneumonia ( $p < 0.05$ ) in analyses restricted to the first year of life (e.g., OR = 4.06; 95% CI: 1.93, 8.57 for a 5-μg/m<sup>3</sup> increase in PM<sub>2.5</sub>) (Table 3).


Further, the combined effect estimate for all associations (pneumonia, otitis media, and croup) increased when analyses were restricted to outcomes in the first year of life.

Stratified meta-analyses suggested slightly stronger effects in females and in those from middle socioeconomic groups (see Supplemental Material, Tables S5 and S6). In analyses stratified by residential mobility during follow-up, the associations between air pollution and respiratory infection were not consistent by strata: Pneumonia effects were greater for movers (OR = 1.62; 95% CI: 1.20, 2.18 vs. 1.21; 95% CI: 0.88, 1.67 for NO<sub>2</sub>), whereas otitis media effects were greater for nonmovers (OR = 1.08; 95% CI: 1.01, 1.16 vs. 1.03; 95% CI: 0.71, 1.48 for NO<sub>2</sub>; see Supplemental Material, Table S7).

Inclusion of additional covariates into the individual cohort models (birth weight, maternal age and area level socioeconomic indicators) did not change air pollution effect estimates or improve model fit (data not shown). There was no consistent evidence for spatial clustering when area-level variables were included as a random effect in models (data not shown). Finally, analyses using back-extrapolated monitoring data were generally consistent with the main findings (see Supplemental Material, Table S8).

## Discussion

As part of the ESCAPE project we had the unprecedented opportunity to examine outdoor air pollution as a risk factor for respiratory infection during early childhood in



**Figure 1.** Forest plots of individual cohort and combined effect estimates (ORs) by outcome for (A) NO<sub>2</sub>, (B) NO<sub>x</sub>, (C) PM<sub>2.5</sub>, (D) PM<sub>2.5</sub> absorbance, (E) PM<sub>10</sub>, (F) coarse PM, (G) traffic intensity on nearest street, and (H) traffic load on all major roads. Weight indicates relative weight (%) assigned using random-effects meta-analysis. The lifetime cumulative incidence of respiratory infection (pneumonia, otitis media, croup) was assessed at 12 months (INMA Valencia), 14 months (INMA Gipuzkoa, INMA Sabadell), 15 months (GASPII), 18 months (INMA Asturias), 24 months (BAMSE, GINI/LISA North, GINI/LISA South, PIAMA), and 36 months (MAAS) of age. Individual cohort models were adjusted for municipality (BAMSE), sex, older siblings, breastfeeding at 6 months, atopy of either parent, any child-care reported during follow-up, maternal smoking during pregnancy, any environmental tobacco smoke in the child's home reported during follow-up, visible mold or dampness in the home, use of gas stove, birth season, parental socioeconomic status (low, medium, high), and intervention (GINIplus, MAAS, PIAMA). Associations are presented for the following increments in exposure: 10 μg/m<sup>3</sup> for NO<sub>2</sub>, 20 μg/m<sup>3</sup> for NO<sub>x</sub>, 5 μg/m<sup>3</sup> for PM<sub>2.5</sub>, 1 unit for PM<sub>2.5</sub> absorbance, 10 μg/m<sup>3</sup> for PM<sub>10</sub>, 5 μg/m<sup>3</sup> for coarse PM, 5,000 vehicles/day for traffic intensity on the nearest street; and 4,000 vehicle-km/day for traffic load on major roads within a 100-m buffer.

an analysis combining 10 European birth cohorts ( $N_{\text{Total}} = 16,059$ ) with data on parent-reported physician-diagnosed pneumonia, otitis media, and croup; we also examined individual air pollution exposure estimates based on common ESCAPE protocols. We found consistent evidence for an association between air pollution and pneumonia, and some evidence for otitis media, during the first two years of life.

Urban air pollution has been associated with respiratory tract infections (Jedrychowski et al. 2013; Lin et al. 2005), pneumonia (Gouveia and Fletcher 2000), croup (Schwartz et al. 1991), persistent cough (Esplugues et al. 2011), and otitis media (MacIntyre et al. 2011) during childhood. Associations have also been reported for indoor air pollution and pneumonia in developing countries (da Costa et al. 2004; Mahalanabis et al. 2002) where concentrations are considerably higher than in our study areas. Our findings are consistent with previous studies that used similar methods to examine air pollution and otitis media in three of our cohorts—PIAMA (Brauer et al. 2006), LISApplus Munich (Brauer et al. 2006), and INMA (Aguilera et al. 2013)—and a recent meta-analysis on long-term  $\text{PM}_{2.5}$  and acute lower respiratory infection in children, which also included the PIAMA study (Mehta et al. 2013).

Similar to secondhand smoke (U.S. Department of Health and Human Services 2006), air pollution is thought to increase susceptibility to respiratory infections primarily via an inflammatory response (Li et al. 2008). Urban air pollution may impair defense mechanisms (Clarke et al. 2000; Leonardi et al. 2000), and oxidant pollutants, in particular, may exacerbate virus-induced inflammation of the respiratory system (Lambert et al. 2003; Spannhake et al. 2002).

Analyses were restricted to the first years of life to include the period of greatest age-specific incidence of respiratory infections (Schnabel et al. 2009; Walker et al. 2013). Our findings suggested that air pollution effects may be slightly stronger during the first year (Table 3). This finding could highlight a unique period of susceptibility when children are at increased risk of respiratory infections due to air pollution (Gehring et al. 2002; Gouveia and Fletcher 2000; Heinrich and Slama 2007). It is also possible that the null findings for infections during the second year of life are due to increased exposure misclassification as older children may spend less time at their home address due to increased child-care enrollment.

A unique strength of LUR models is their ability to capture small-scale spatial variability in exposure; however, the measurements

used to create the ESCAPE exposure models were taken after the birth year (Eeftens et al. 2012a; Cyrys et al. 2012), and this may have introduced exposure misclassification. Although it is possible that overall levels of air pollution changed during this period, previous findings suggest that the spatial distribution of air pollutants within each area remained consistent (Cesaroni et al. 2012; Eeftens et al. 2011; Wang et al. 2013). Further, our sensitivity analyses using monitoring data to back-extrapolate exposure estimates to the actual first year of life were consistent with our main findings (see Supplemental Material, Table S8).

The wording of parental questionnaires was similar across each cohort, and previous research has shown good agreement between maternal recall and medical records during early childhood (D'Souza-Vazirani et al. 2005; Vernacchio et al. 2007). Geographic differences in the prevalence of outcomes across the cohorts were most pronounced for otitis media and may point to potential diagnostic biases or disease misclassification between countries. It was not possible to adjust for epidemics, the impact of vaccinations, or the frequency of infections because data were not available across all cohorts. Furthermore, defining upper respiratory tract infections (otitis media, croup) by physician




Figure 1. Continued.

**Table 2.** Combined estimates from random-effects meta-analyses for residential air pollution and respiratory infections during early life (up to 36 months).<sup>a</sup>

	Crude <sup>b</sup>			Adjusted <sup>c</sup>		
	OR (95% CI)	I <sup>2</sup>	p-Value	OR (95% CI)	I <sup>2</sup>	p-Value
Pneumonia						
NO <sub>2</sub>	1.25 (1.04, 1.50)*	37.1	0.112	1.30 (1.02, 1.65)*	52.9	0.024
NO <sub>x</sub>	1.23 (1.06, 1.41)*	22.2	0.239	1.26 (1.04, 1.52)*	44.0	0.066
PM <sub>2.5</sub>	2.13 (0.82, 5.49)	79.7	0.000	2.58 (0.91, 7.27)	81.7	0.000
PM <sub>2.5</sub> absorbance	1.78 (1.30, 2.43)*	0	0.734	1.99 (1.44, 2.75)*	0	0.663
PM <sub>10</sub>	1.55 (1.03, 2.34)*	29.2	0.205	1.76 (1.00, 3.09)*	51.2	0.051
Coarse PM	1.23 (1.02, 1.47)*	0	0.626	1.24 (1.03, 1.50)*	0	0.579
Traffic, nearest street	1.08 (1.03, 1.14)*	0	0.997	1.09 (1.03, 1.15)*	0	0.969
Traffic, major streets	1.19 (1.08, 1.31)*	0	0.979	1.21 (1.09, 1.34)*	0	0.843
Otitis media						
NO <sub>2</sub>	1.08* (1.01, 1.15)	4.8	0.395	1.09 (1.02, 1.16)*	0	0.515
NO <sub>x</sub>	1.04 (0.98, 1.10)	0.5	0.430	1.05 (0.98, 1.12)	0	0.458
PM <sub>2.5</sub>	1.02 (0.71, 1.45)	55.5	0.047	1.06 (0.75, 1.49)	47.9	0.088
PM <sub>2.5</sub> absorbance	1.05 (0.80, 1.37)	46.7	0.095	1.08 (0.83, 1.39)	39.9	0.139
PM <sub>10</sub>	0.98 (0.83, 1.17)	11.6	0.341	0.98 (0.84, 1.14)	0	0.539
Coarse PM	0.96 (0.87, 1.06)	0	0.608	0.97 (0.88, 1.08)	0	0.805
Traffic, nearest street	0.98 (0.94, 1.03)	1.4	0.385	0.98 (0.93, 1.02)	0	0.497
Traffic, major streets	1.00 (0.91, 1.09)	0	0.462	0.99 (0.89, 1.10)	18.2	0.300
Croup						
NO <sub>2</sub>	0.92 (0.80, 1.07)	0	0.884	0.96 (0.83, 1.12)	0	0.909
NO <sub>x</sub>	0.96 (0.83, 1.10)	0	0.895	0.99 (0.86, 1.14)	0	0.936
PM <sub>2.5</sub>	0.83 (0.58, 1.19)	0	0.760	0.90 (0.63, 1.30)	0	0.703
PM <sub>2.5</sub> absorbance	0.95 (0.66, 1.37)	5.0	0.368	1.03 (0.72, 1.47)	0	0.554
PM <sub>10</sub>	0.89 (0.70, 1.13)	0	0.586	0.92 (0.72, 1.18)	0	0.595
Coarse PM	0.95 (0.80, 1.12)	0	0.551	0.97 (0.82, 1.15)	0	0.787
Traffic, nearest street	0.98 (0.93, 1.04)	0	0.926	0.99 (0.93, 1.05)	0	0.853
Traffic, major streets	0.97 (0.86, 1.09)	0	0.734	0.98 (0.87, 1.11)	0	0.901

Associations are presented for the following increments in exposure: 10 µg/m<sup>3</sup> for NO<sub>2</sub>, 20 µg/m<sup>3</sup> for NO<sub>x</sub>, 5 µg/m<sup>3</sup> for PM<sub>2.5</sub>, 1 unit for PM<sub>2.5</sub> absorbance, 10 µg/m<sup>3</sup> for PM<sub>10</sub>, 5 µg/m<sup>3</sup> for coarse PM, 5,000 vehicles/day for traffic intensity on the nearest street; and 4,000 vehicle-km/day for traffic load on major roads within a 100-m buffer; associations with traffic intensity and traffic load were additionally adjusted for background NO<sub>2</sub> concentrations.

<sup>a</sup>Outcomes assessed up to 12 months (INMA Valencia), 14 months (INMA Gipuzkoa, INMA Sabadell), 15 months (GASPII), 18 months (INMA Asturias), 24 months (BAMSE, GINIplus, LISAPLUS, PIAMA), and 36 months (MAAS). <sup>b</sup>Crude models were adjusted for sex and municipality (BAMSE). <sup>c</sup>Adjusted models included municipality (BAMSE), sex, older siblings, breastfeeding at 6 months, atopy of either parent, any child-care reported during follow-up, maternal smoking during pregnancy, any environmental tobacco smoke in the child's home reported during follow-up, visible mold or dampness in the home, use of gas stove, birth season, parental socioeconomic status (low, medium, high), and intervention (GINIplus, MAAS, PIAMA). \**p* < 0.05.

**Table 3.** Adjusted combined estimates for air pollution exposure at the birth address and respiratory infection by year of life [OR (95% CI)].

	Pneumonia <sup>a</sup> ( <i>n</i> = 12,891)	Otitis media <sup>b</sup> ( <i>n</i> = 8,722)	Croup <sup>c</sup> ( <i>n</i> = 9,101)
Respiratory infections during the first year <sup>d</sup> of life			
NO <sub>2</sub>	1.47* (1.15, 1.89)	1.19* (1.07, 1.33)	1.05 (0.83, 1.32)
NO <sub>x</sub>	1.45* (1.21, 1.75)	1.09 (0.98, 1.22)	1.10 (0.90, 1.36)
PM <sub>2.5</sub>	4.06* (1.93, 8.57)	1.21 (0.64, 2.28)	1.15 (0.67, 1.97)
PM <sub>2.5</sub> absorbance	2.71* (1.68, 4.37)	1.32 (0.99, 1.75)	1.04 (0.59, 1.83)
PM <sub>10</sub>	1.77* (1.18, 2.67)	1.24 (0.76, 2.02)	1.07 (0.75, 1.53)
Coarse PM	1.46* (1.11, 1.92)	1.16 (0.80, 1.70)	1.02 (0.80, 1.30)
Traffic, nearest street	1.14* (1.07, 1.22)	0.99 (0.94, 1.04)	1.03 (0.94, 1.13)
Traffic, major streets	1.31* (1.15, 1.50)	1.03 (0.93, 1.14)	1.00 (0.81, 1.24)
Respiratory infections during the second year <sup>e</sup> of life			
NO <sub>2</sub>	1.40* (1.04, 1.88)	1.07 (0.96, 1.20)	0.92 (0.78, 1.09)
NO <sub>x</sub>	1.29* (1.07, 1.55)	1.02 (0.89, 1.17)	0.92 (0.78, 1.08)
PM <sub>2.5</sub>	2.65 (0.63, 11.2)	1.06 (0.64, 1.74)	0.76 (0.51, 1.15)
PM <sub>2.5</sub> absorbance	1.90 (0.93, 3.87)	1.20 (0.80, 1.79)	0.89 (0.59, 1.35)
PM <sub>10</sub>	1.42 (0.99, 2.03)	1.00 (0.84, 1.19)	0.83 (0.63, 1.09)
Coarse PM	1.24 (0.98, 1.56)	1.00 (0.89, 1.13)	0.89 (0.73, 1.08)
Traffic, nearest street	1.05 (0.98, 1.13)	0.96 (0.90, 1.03)	0.93 (0.81, 1.07)
Traffic, major streets	1.10 (0.90, 1.34)	0.96 (0.83, 1.10)	1.00 (0.88, 1.14)

Associations are presented for the following increments in exposure: 10 µg/m<sup>3</sup> for NO<sub>2</sub>, 20 µg/m<sup>3</sup> for NO<sub>x</sub>, 5 µg/m<sup>3</sup> for PM<sub>2.5</sub>, 1 unit for PM<sub>2.5</sub> absorbance, 10 µg/m<sup>3</sup> for PM<sub>10</sub>, 5 µg/m<sup>3</sup> for coarse PM, 5,000 vehicles/day for traffic intensity on the nearest street; and 4,000 vehicle-km/day for traffic load on major roads within a 100-m buffer; associations with traffic intensity and traffic load were additionally adjusted for background NO<sub>2</sub> concentrations.

<sup>a</sup>Based on four studies: BAMSE, GINI/LISA North, GINI/LISA South, PIAMA. <sup>b</sup>Based on 3 studies: BAMSE, LISAPLUS North, LISAPLUS South, PIAMA. <sup>c</sup>Based on three studies: BAMSE, GINI/LISA North, GINI/LISA South. <sup>d</sup>Defined as 0–12 months. <sup>e</sup>Defined as 13–24 months. Models were adjusted for municipality (BAMSE), sex, older siblings, breastfeeding at 6 months, atopy of either parent, any child-care reported during follow-up, maternal smoking during pregnancy, any environmental tobacco smoke in the child's home reported during follow-up, visible mold or dampness in the home, use of gas stove, birth season, parental socioeconomic status (low, medium, high), and intervention (GINIplus, PIAMA). \**p* < 0.05.

diagnosis is complicated by the fact that not all infections present with acute symptoms severe enough to warrant a physician visit, in contrast with pneumonia, which routinely presents with a high fever and/or difficulty breathing (Edmond et al. 2012).

## Conclusion

Our meta-analysis of 10 European birth cohorts found consistent evidence for an association between traffic-related air pollution and pneumonia, and some evidence to suggest an association with otitis media. Policies aimed at reducing air pollution may be successful in reducing the overall burden of pneumonia in early childhood.

## CORRECTION

The value for “Traffic load on major streets within 100-m buffer” for GINI/LISA South in Table 1 was incorrect in the manuscript originally published online. It has been corrected here.

## REFERENCES

- Aguilera I, Pedersen M, Garcia-Esteban R, Ballester F, Basterrechea M, Esplugues A, et al. 2013. Early-life exposure to outdoor air pollution and respiratory health, ear infections, and eczema in infants from the INMA study. *Environ Health Perspect* 121:387–392; doi:10.1289/ehp.1205281.
- Bateson TF, Schwartz J. 2008. Children's response to air pollutants. *J Toxicol Environ Health A* 71(3):238–243.
- Beelen R, Hoek G, Vienneau D, Eeftens M, Dimakopoulou M, Pedeli X, et al. 2013. Development of NO<sub>2</sub> and NO<sub>x</sub> land use regression models for estimating air pollution exposure in 36 study areas in Europe—the ESCAPE project. *Atmos Environ* 72:10–23.
- Black RE, Cousens S, Johnson HL, Lawn JE, Rudan I, Bassani DG, et al. 2010. Global, regional, and national causes of child mortality in 2008: a systematic analysis. *Lancet* 375:1969–1987.
- Brauer M, Gehring U, Brunekreef B, de Jongste J, Gerritsen J, Rovers M, et al. 2006. Traffic-related air pollution and otitis media. *Environ Health Perspect* 114:1414–1418; doi:10.1289/ehp.9089.
- Brunekreef B, Smit J, de Jongste J, Neijens H, Gerritsen J, Postma D, et al. 2002. The Prevention and Incidence of Asthma and Mite Allergy (PIAMA) birth cohort study: design and first results. *Pediatr Allergy Immunol* 13:55–60.
- Cesaroni G, Porta D, Badaloni C, Stafoggia M, Eeftens M, Meliefste K, et al. 2012. Nitrogen dioxide levels estimated from land use regression models several years apart and association with mortality in a large cohort study. *Environ Health* 11:48; doi:10.1186/1476-069X-11-48.
- Clarke RW, Antonini JM, Hemenway DR, Frank R, Kleeberger SR, Jakab GJ. 2000. Inhaled particle-bound sulfate: effects on pulmonary inflammatory responses and alveolar macrophage function. *Inhal Toxicol* 12(3):169–186.
- Custovic A, Simpson BM, Murray CS, Lowe L, Woodcock A; NAC Manchester Asthma and Allergy Study Group. 2002. The National Asthma Campaign Manchester Asthma and Allergy Study. *Pediatr Allergy Immunol* 13(suppl 15):32–37.
- Cyrys J, Eeftens M, Heinrich J, Ampe C, Armengaud A, Beelen R, et al. 2012. Variation of NO<sub>2</sub> and NO<sub>x</sub> concentrations between and within 36 European study areas: results from the ESCAPE study. *Atmos Environ* 62: 374–390.
- da Costa JL, Navarro A, Neves JB, Martin M. 2004. Household wood and charcoal smoke increases risk of otitis media in childhood in Maputo. *Int J Epidemiol* 33(3):573–578.
- DerSimonian R, Laird N. 1986. Meta-analysis in clinical trials. *Control Clin Trials* 7(3):177–188.
- D'Souza-Vazirani D, Minkovitz CS, Strobino DM. 2005. Validity



- of maternal report of acute health care use for children younger than 3 years. *Arch Pediatr Adolesc Med* 159(2):167–172.
- Edmond K, Scott S, Korczak V, Ward C, Sanderson C, Theodoratou E, et al. 2012. Long term sequelae from childhood pneumonia; systematic review and meta-analysis. *PLoS One* 7(2):e31239; doi:10.1371/journal.pone.0031239.
- Eeftens M, Beelen R, de Hoogh K, Bellander T, Cesaroni G, Cirach M, et al. 2012a. Development of land use regression models for PM<sub>2.5</sub>, PM<sub>2.5</sub> absorbance, PM<sub>10</sub> and PMcoarse in 20 European study areas; results of the ESCAPE project. *Environ Sci Technol* 46(20):11195–11205.
- Eeftens M, Beelen R, Fischer P, Brunekreef B, Meliefste K, Hoek G. 2011. Stability of measured and modelled spatial contrasts in NO<sub>2</sub> over time. *Occup Environ Med* 68(10):765–770.
- Eeftens M, Tsai M-Y, Ampe C, Anwander B, Beelen R, Bellander T, et al. 2012b. Spatial variation of PM<sub>2.5</sub>, PM<sub>10</sub>, PM<sub>2.5</sub> absorbance and PMcoarse concentrations between and within 20 European study areas and the relationship with NO<sub>2</sub>—Results of the ESCAPE project. *Atmos Environ* 62(0):303–317.
- ESCAPE (European Study of Cohorts for Air Pollution Effects). 2009. Home page. Available: <http://www.escapeproject.eu/> [accessed 22 November 2013].
- ESCAPE (European Study of Cohorts for Air Pollution Effects). 2013. Manuals. Available: <http://www.escapeproject.eu/manuals/> [accessed 22 November 2013].
- Espugues A, Ballester F, Estarlich M, Llop S, Fuentes-Leonarte V, Mantilla E, et al. 2011. Outdoor, but not indoor, nitrogen dioxide exposure is associated with persistent cough during the first year of life. *Sci Total Environ* 409(22):4667–4673.
- Gehring U, Cyrys J, Sedlmeir G, Brunekreef B, Bellander T, Fischer P, et al. 2002. Traffic-related air pollution and respiratory health during the first 2 yrs of life. *Eur Respir J* 19(4):690–698.
- Gouveia N, Fletcher T. 2000. Respiratory diseases in children and outdoor air pollution in São Paulo, Brazil: a time series analysis. *Occup Environ Med* 57(7):477–483.
- Guxens M, Aguilera I, Ballester F, Estarlich M, Fernández-Somoano A, Lertxundi A, et al. 2012. Prenatal exposure to residential air pollution and infant mental development: modulation by antioxidants and detoxification factors. *Environ Health Perspect* 120:144–149; doi:10.1289/ehp.1103469.
- Heinrich J, Bolte G, Hölischer B, Douwes J, Lehmann I, Fahlbusch B, et al. 2002. Allergens and endotoxin on mothers' mattresses and total immunoglobulin E in cord blood of neonates. *Eur Respir J* 20(3):617–623.
- Heinrich J, Slama R. 2007. Fine particles, a major threat to children. *Int J Hyg Environ Health* 210(5):617–22.
- Huedo-Medina TB, Sánchez-Meca J, Marín-Martínez F, Botella J. 2006. Assessing heterogeneity in meta-analysis: Q statistic or  $I^2$  index? *Psychol Methods* 11(2):193–206.
- Jedrychowski WA, Perera FP, Spengler JD, Mroz E, Stigter L, Flak E, et al. 2013. Intrauterine exposure to fine particulate matter as a risk factor for increased susceptibility to acute broncho-pulmonary infections in early childhood. *Int J Hyg Environ Health* 216(4):395–401.
- Lambert AL, Mangum JB, DeLorme MP, Everitt JI. 2003. Ultrafine carbon black particles enhance respiratory syncytial virus-induced airway reactivity, pulmonary inflammation, and chemokine expression. *Toxicol Sci* 72(2):339–346.
- Leonardi GS, Houthuijs D, Steerenberg PA, Fletcher T, Armstrong B, Antova T, et al. 2000. Immune biomarkers in relation to exposure to particulate matter: a cross-sectional survey in 17 cities of Central Europe. *Inhal Toxicol* 12(suppl 4):1–14.
- Li N, Xia T, Nel AE. 2008. The role of oxidative stress in ambient particulate matter-induced lung diseases and its implications in the toxicity of engineered nanoparticles. *Free Radic Biol Med* 44(9):1689–1699.
- Lin M, Stieb DM, Chen Y. 2005. Coarse particulate matter and hospitalization for respiratory infections in children younger than 15 years in Toronto: a case-cross-over analysis. *Pediatrics* 116(2):e235–e240.
- MacIntyre EA, Karr CJ, Koehoorn M, Demers PA, Tamburic L, Lencar C, et al. 2011. Residential air pollution and otitis media during the first two years of life. *Epidemiology* 22(1):81–89.
- Mahalanabis D, Gupta S, Paul D, Gupta A, Lahiri M, Khaled MA. 2002. Risk factors for pneumonia in infants and young children and the role of solid fuel for cooking: a case-control study. *Epidemiol Infect* 129(1):65–71.
- Mehta S, Shin H, Burnett R, North T, Cohen AJ. 2013. Ambient particulate air pollution and acute lower respiratory infections: a systematic review and implications for estimating the global burden of disease. *Air Qual Atmos Health* 6(1):69–83.
- Porta D, Forastiere F, Di Lallo D, Perucci CA; Gruppo Collaborativo GASPII. 2007. Enrolment and follow-up of a birth cohort in Rome [in Italian]. *Epidemiol Prev* 31(6):303–308.
- Schnabel E, Sausenthaler S, Brockow I, Liese J, Herbarth O, Michael B, et al. 2009. Burden of otitis media and pneumonia in children up to 6 years of age: results of the LISA birth cohort. *Eur J Pediatr* 168(10):1251–1257.
- Schwartz J, Spix C, Wichmann HE, Malin E. 1991. Air pollution and acute respiratory illness in five German communities. *Environ Res* 56:1–14.
- Spannhake EW, Reddy SP, Jacoby DB, Yu XY, Saatian B, Tian J. 2002. Synergism between rhinovirus infection and oxidant pollutant exposure enhances airway epithelial cell cytokine production. *Environ Health Perspect* 110:665–670.
- U.S. Department of Health and Human Services. 2006. The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General. Washington, DC:U.S. Department of Health and Human Services.
- Vernacchio L, Vezina RM, Ozonoff A, Mitchell AA. 2007. Validity of parental reporting of recent episodes of acute otitis media: a Slone Center Office-Based Research (SCOR) Network study. *J Am Board Fam Med* 20(2):160–163.
- Walker CLF, Rudan I, Liu L, Nair H, Theodoratou E, Bhutta ZA, et al. 2013. Global burden of childhood pneumonia and diarrhoea. *Lancet* 381:1405–1416.
- Wang R, Henderson SB, Sbihi H, Allen RW, Brauer M. 2013. Temporal stability of land use regression models for traffic-related air pollution. *Atmos Environ* 64:312–319.
- Wickman M, Kull I, Pershagen G, Nordvall SL. 2002. The BAMSE project: presentation of a prospective longitudinal birth cohort study. *Pediatr Allergy Immunol* 13(suppl 15):11–13.
- Williams BG, Gouws E, Boschi-Pinto C, Bryce J, Dye C. 2002. Estimates of world-wide distribution of child deaths from acute respiratory infections. *Lancet Infect Dis* 2:25–32.
- Zirngibl A, Franke K, Gehring U, von Berg A, Berdel D, Bauer CP, et al. 2002. Exposure to pets and atopic dermatitis during the first two years of life. A cohort study. *Pediatr Allergy Immunol* 13(6):394–401.